

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/54, 1/15, C12Q 1/68 C12N 5/10, 9/10	A2	(11) International Publication Number: WO 93/13207 (43) International Publication Date: 8 July 1993 (08.07.93)
(21) International Application Number: PCT/US92/11353 (22) International Filing Date: 30 December 1992 (30.12.92) (30) Priority data: 07/816,284 31 December 1991 (31.12.91) US (71) Applicant: ZYMOGENETICS, INC. [US/US]; 4225 Roosevelt Way N.E., Seattle, WA 98105 (US). (72) Inventors: O'HARA, Patrick, J. ; 515 North 64th, Seattle, WA 98103 (US). GRANT, Francis, J. ; 7714 - 37th Ave- nue N.E., Seattle, WA 98115 (US). SHEPPARD, Paul, O. ; 20717 N.E. 2nd, Redmond, WA 98199 (US).		(74) Agents: PARMELEE, Steven, W. et al.; Townsend and Townsend, One Market Plaza, 20th Fl., Steuart Tower, San Francisco, CA 94105 (US). (81) Designated States: AU, CA, FI, JP, NO, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NOVEL HUMAN TRANSGLUTAMINASES (57) Abstract Human prostatic and placental transglutaminases are identified and cloned. The human transglutaminases herein are use- ful for, <i>inter alia</i> , therapeutic wound repair, closure of skin grafts, stabilizing food preparations, and markers for identifying agents which act as agonists or antagonists of cellular apoptosis.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

5

NOVEL HUMAN TRANSGLUTAMINASESBackground of the Invention

10 Transglutaminases are a group of calcium dependent
enzymes that catalyze the crosslinking of proteins by
promoting the formation of ϵ -(γ -glutaminyll)lysine isopeptide
bonds between protein-bound glutamine and lysine residues.
These enzymes are believed to be widely distributed in nature,
15 as the crosslinks are found in both prokaryotic and eukaryotic
cells. Although different transglutaminases appear to be very
similar in substrate specificity, several distinct forms of
the enzymes have been identified. See generally, Folk, Ann.
Rev. Biochem. 49:517-531 (1980).

20 Transglutaminase-mediated protein crosslinking
reactions have been implicated in both normal and pathological
processes in mammalian cells and tissues. The crosslink may
act to maintain some forms of protein structure, such as in
the terminal differentiation of epidermal cell layers and in
25 other cellular architecture. An intracellular
transglutaminase known as epidermal or Type I transglutaminase
has been isolated and cloned from rabbit epithelial cells
(Floyd and Jetten, Mol. Cell. Biol. 9:4846-4851 (1989)), and a
transglutaminase has been isolated and cloned from guinea pig
30 liver cells (Ikura et al., Biochem. 27: 2898-2905 (1988)).
Other transglutaminase activities have been described
including hair follicle transglutaminase, keratinocyte
transglutaminase, and prostate transglutaminase (Wilson et
al., Fed. Proc. 38:1809 (1979)). Lee et al., Prep. Biochem.
35 16:321-335 (1986) have described the purification of a
transglutaminase from human erythrocytes. These

transglutaminases have been shown to be distinct from a plasma transglutaminase, Factor XIII, an enzyme whose primary function appears to be stabilizing fibrin clots. Factor XIII has also been purified, cloned, and sequenced. (Ichinose, et al., Biochem. 25:6900-6906 (1986), Takahashi, et al., Proc. Natl. Acad. Sci. U.S.A. 83:8018-8023 (1986)).

Transglutaminases have been employed for crosslinking purposes in a variety of fields. Certain microbial transglutaminases have found use in food technology to add texture to processed foods, particularly fish and cheese. Others have been used in enzyme-catalyzed fluorescent labeling of proteins, in the introduction of cleavable crosslinks, and in the solid-phase reversible removal of specific proteins from biological systems. Factor XIII preparations have been proposed for a variety of therapeutic uses, such as the treatment of subarachnoid hemorrhage and inflammatory bowel disease.

Presently, a plasma derived Factor XIII is available as a fibrin sealant, but, as with most plasma-derived products, carries an inherent risk of viral contamination. Further, Factor XIII and certain other transglutaminases are zymogens, requiring some form of activation to become catalytically active. And, as each transglutaminase has a restricted range of substrates, their activity may be limited in certain applications. Accordingly, what is needed in the art are methods for producing by recombinant means human transglutaminases. The present invention fulfills these and other related needs.

30

Summary of the Invention

The present invention provides the ability to produce human prostatic and placental transglutaminases and polypeptides or fragments thereof by recombinant means,

35

preferably in cultured eukaryotic cells. The expressed transglutaminase may or may not have the biological activity of the native enzyme, depending on the intended use.

Accordingly, isolated and purified polynucleotides are

described which code for the transglutaminases and fragments thereof, where the polynucleotides may be in the form of DNA, such as cDNA or genomic DNA, or RNA. Based on these sequences probes may be designed for hybridization to identify these and related genes or transcription products thereof which encode human prostatic and placental transglutaminases.

In related embodiments the invention concerns DNA constructs which comprise a transcriptional promoter, a DNA sequence which encodes the prostatic or placental transglutaminase or fragment thereof, and a transcriptional terminator, each operably linked for expression of the enzyme or enzyme fragment. The constructs are preferably used to transform or transfect host cells, preferably eukaryotic cells, more preferably yeast or mammalian cells. For large scale production the expressed prostatic or placental transglutaminase may be isolated from the cells by, for example, immunoaffinity purification.

Nucleic acid sequences which encode the human prostatic or placental transglutaminases of the invention and the recombinant transglutaminases themselves can also be used to develop compounds which can alter transglutaminase-associated apoptosis of a eukaryotic cell. Compounds may be screened for agonistic or antagonistic effects on transglutaminase-mediated metabolism in the host cell.

Description of the Specific Embodiments

The present invention provides isolated polynucleotide molecules encoding human prostatic and placental transglutaminases, thereby providing for the expression of human prostatic and placental transglutaminase polypeptides and fragments thereof. Isolated polynucleotide molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are provided free of other genes with which they are naturally associated and may include naturally occurring 5' and 3' untranslated sequences that represent regulatory regions such as promoters and terminators. The identification of regulatory regions within the naturally occurring 5' and 3' untranslated regions will be evident to one of ordinary skill in the art (for review, see Dynan and Tijan, Nature 316: 774-778, 1985; Birnstiel et al., Cell 41: 349-359, 1985; Proudfoot, Trends in Biochem. Sci. 14: 105-110, 1989; and Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY, 1989; which are incorporated herein by reference).

As will be understood by one skilled in the art, the DNA molecules of the present invention encompass allelic variants and genetically engineered or synthetic variants of the transglutaminases that encode conservative amino acid substitutions and/or minor additions, or deletions of amino acids. Such variants also encompass DNA molecules containing degeneracies in the DNA code wherein host-preferred codons are substituted for the analogous codons in the human sequence. In addition, substantially similar DNA molecules of the present invention encompass those DNA molecules that are capable of hybridizing to the DNA sequences of the present invention under high or low stringency (see Sambrook et al., *ibid.*) and those sequences that are degenerate as a result of

the genetic code to the amino acid sequences of the present invention.

Recombinant DNA expression systems provide convenient means for obtaining large quantities of the human transglutaminases in relatively pure form. By human prostatic or placental transglutaminase polypeptides and fragments is meant to include sequences of amino acids from 9 to 20 amino acids up to entire proteins, which have at least about 85% homology, preferably at least 90%, and more preferably at least about 95% or more homology to the amino acid sequences of the human prostatic or placental transglutaminases of the invention. As will be appreciated by those skilled in the art, the invention also includes those polypeptides having slight variations in amino acid sequences or other properties. Such variations may arise naturally as allelic variations (e.g., due to genetic polymorphism) or may be produced by human intervention (e.g., by mutagenesis of cloned DNA sequences), such as induced point, deletion and insertion mutations.

Nucleic acid molecules encoding the human transglutaminases as described herein can be cloned from a variety of human cell sources that express the enzymes. Preferred sources for human prostatic transglutaminase include human prostate or liver cells and tissues, and for human placental transglutaminase include, e.g., human placental tissue. Useful isolated nucleic acid sequences of the invention which encode the human transglutaminases include mRNA, genomic DNA and cDNA. For expression, cDNAs are generally preferred because they lack introns that may interfere with expression.

To obtain human prostate and/or placental transglutaminase clones, a human prostate tissue cDNA library and/or human placental tissue cDNA library is amplified to obtain DNA molecules encoding transglutaminases using oligonucleotide primers in a polymerase chain reaction ("PCR";

U.S. Patent Nos. 4,683,195, 4,683,202, incorporated herein by reference). The oligonucleotide primer sequences are designed by preparing a multiple sequence alignment of sequence information for a variety of transglutaminases and related proteins (e.g., rat keratinocyte transglutaminase, human keratinocyte transglutaminase, human transglutaminase K, human factor XIII, human endothelial cell transglutaminase, mouse macrophage transglutaminase, guinea pig transglutaminase, human erythrocyte membrane protein band 4.2, rabbit transglutaminase type I, and bovine factor XIII). The multiple alignment is subjected to analysis for least degenerate/most conserved regions from which primers, which are generally about 17-20 bases long, are designed. Primers were designed from three regions of multiple homology in Example I described below: one from the active site region, and two from other regions which seemed to have structural importance, based on, inter alia, the presence of hydrophobic residues and proline residues. Following amplification and enrichment for the desired DNA molecules, the molecules are identified and used to screen and isolate full length cDNA clones for the prostate and placental transglutaminases.

cDNA libraries can be screened with, e.g., labeled probes from random-primed DNA molecules encoding human prostatic or placental transglutaminase, which probes preferably span the enzyme's active site and/or putative calcium binding site. To obtain the human placental transglutaminase clone, an oligo-d(T) primed cDNA library can be constructed from poly(A)⁺ RNA purified from human placental tissues. Partial clones may be used as probes in additional screening until the complete coding sequence is obtained.

In addition to the use of partial clones to obtain full length transglutaminase clones, PCR amplification may be used to obtain a complete cDNA. Synthetic oligonucleotide primers may be designed to hybridize to vector sequences near the cDNA insert boundary and to DNA sequences within the

transglutaminase coding sequence. Polymerase chain amplification may be used in conjunction with such primers to obtain DNA segments encoding terminal DNA sequences for completing a partial cDNA clone.

5 If necessary, partial clones are joined in the correct reading frame to construct the complete coding sequence. Joining is achieved by, for example, digesting clones with appropriate restriction endonucleases and joining the fragments enzymatically in the proper orientation.

10 Depending on the fragments and the particular restriction endonucleases chosen, it may be necessary to remove unwanted DNA sequences through a "loop out" process of deletion mutagenesis or through a combination of restriction endonuclease cleavage and mutagenesis. It is preferred that

15 the resultant sequence be in the form of a continuous open reading frame, that is, that it lack intervening sequences (introns). The sequence of one exemplary human prostate clone described herein is shown in SEQ. ID. NO. 14.

 With the nucleotide and deduced amino acid

20 sequences of human prostate transglutaminase provided herein, genomic or cDNA sequences encoding human prostatic transglutaminase may be obtained from libraries prepared from other cells and tissues according to known procedures. For instance, using oligonucleotide probes derived from human

25 prostate transglutaminase sequences, generally of at least about fourteen nucleotides and up to twenty-five or more nucleotides in length, DNA sequences encoding transglutaminases of other cells or tissues may be obtained. If partial clones are obtained, it is necessary to join them

30 in proper reading frame to produce a full length clone, using such techniques as endonuclease cleavage, ligation and loopout mutagenesis.

 For expression, a DNA sequence encoding human prostate or placental transglutaminase polypeptide is inserted

35 into a suitable expression vector, which in turn is used to

transform or transfect appropriate host cells for expression. Expression vectors for use in carrying out the present invention will generally comprise a promoter capable of directing the transcription of a cloned DNA and a transcriptional terminator, operably linked with the sequence encoding the prostate or placental transglutaminase polypeptide so as to produce a continuously transcribable gene sequence which produces sequences in reading frame and is continuously translated to produce a human prostate or placental transglutaminase polypeptide. The expression vectors of the present invention may further include enhancers and other elements such as secretory signal sequences to facilitate expression and/or secretion of the protein. One or more selectable markers may also be included.

Secretory signal sequences, also called leader sequences, prepro sequences and/or pre sequences, are amino acid sequences that act to direct the secretion of mature polypeptides or proteins from a cell. Such sequences are characterized by a core of hydrophobic amino acids and are typically (but not exclusively) found at the amino termini of newly synthesized proteins. Very often the secretory peptide is cleaved from the mature protein during secretion. Such secretory peptides contain processing sites that allow cleavage of the secretory peptides from the mature proteins as they pass through the secretory pathway. A preferred processing site is a dibasic cleavage site, such as that recognized by the Saccharomyces cerevisiae KEX2 gene. A particularly preferred processing site is a Lys-Arg processing site. Processing sites may be encoded within the secretory peptide or may be added to the peptide by, for example, in vitro mutagenesis.

The choice of a suitable secretory signal sequence is well within the level of ordinary skill in the art and will depend on the selected host system employed. Preferred secretory signals include the α factor signal sequence (prepro

sequence: Kurjan and Herskowitz, Cell 30: 933-943, 1982; Kurjan et al., U.S. Patent No. 4,546,082; Brake, U.S. Patent No. 4,870,008), the PHO5 signal sequence (Beck et al., WO 86/00637), the BAR1 secretory signal sequence (MacKay et al., U.S. Patent No. 4,613,572; MacKay, WO 87/002670), the SUC2 signal sequence (Carlsen et al., Molecular and Cellular Biology 3: 439-447, 1983), the α -1-antitrypsin signal sequence (Kurachi et al., Proc. Natl. Acad. Sci. USA 78: 6826-6830, 1981), and the α -2 plasmin inhibitor signal sequence (Tone et al., J. Biochem. (Tokyo) 102:1033-1042, 1987). A particularly preferred signal sequence is the tissue plasminogen activator signal sequence (Pennica et al., Nature 301: 214-221, 1983). Alternately, a secretory signal sequence may be synthesized according to the rules established, for example, by von Heinje (15 European Journal of Biochemistry 133: 17-21, 1983; Journal of Molecular Biology 184: 99-105, 1985; Nucleic Acids Research 14: 4683-4690, 1986).

Secretory signal sequences may be used singly or may be combined. For example, a first secretory signal sequence may be used in combination with a sequence encoding the third domain of barrier (described in U.S. Patent No. 5,037,243, which is incorporated by reference herein in its entirety). The third domain of barrier may be positioned in proper reading frame 3' of the DNA segment of interest or 5' to the DNA segment and in proper reading frame with both the secretory signal sequence and a DNA segment of interest.

Host cells for use in practicing the present invention include mammalian, avian, plant, insect, bacterial and fungal cells, but preferably eukaryotic cells. Preferred eukaryotic cells include cultured mammalian cell lines (e.g., rodent or human cell lines) and fungal cells, including species of yeast (e.g., Saccharomyces spp., particularly S. cerevisiae, Schizosaccharomyces spp., or Kluyveromyces spp.) or filamentous fungi (e.g., Aspergillus spp., Neurospora spp.). Methods for producing recombinant proteins in a

variety of prokaryotic and eukaryotic host cells are generally known in the art.

Suitable yeast vectors for use in the present invention include YRp7 (Struhl et al., Proc. Natl. Acad. Sci. USA 76: 1035-1039, 1978), YEpl3 (Broach et al., Gene 8: 121-133, 1979), POT vectors (Kawasaki et al, U.S. Patent No. 4,931,373, which is incorporated by reference herein), pJDB249 and pJDB219 (Beggs, Nature 275:104-108, 1978) and derivatives thereof. Such vectors will generally include a selectable marker, which may be one of any number of genes that exhibit a dominant phenotype for which a phenotypic assay exists to enable transformants to be selected. Preferred selectable markers are those that complement host cell auxotrophy, provide antibiotic resistance or enable a cell to utilize specific carbon sources, and include LEU2 (Broach et al., ibid.), URA3 (Botstein et al., Gene 8: 17, 1979), HIS3 (Struhl et al., ibid.) or POT1 (Kawasaki et al., ibid.). Another suitable selectable marker is the CAT gene, which confers chloramphenicol resistance on yeast cells.

Preferred promoters for use in yeast include promoters from yeast glycolytic genes (Hitzeman et al., J. Biol. Chem. 255: 12073-12080, 1980; Alber and Kawasaki, J. Mol. Appl. Genet. 1: 419-434, 1982; Kawasaki, U.S. Patent No. 4,599,311) or alcohol dehydrogenase genes (Young et al., in Genetic Engineering of Microorganisms for Chemicals, Hollaender et al., (eds.), p. 355, Plenum, New York, 1982; Ammerer, Meth. Enzymol. 101: 192-201, 1983). In this regard, particularly preferred promoters are the TPI1 promoter (Kawasaki, U.S. Patent No. 4,599,311, 1986) and the ADH2-4^C promoter (Russell et al., Nature 304: 652-654, 1983; Irani and Kilgore, U.S. Patent Application Serial No. 07/631,763 and EP 284,044, which are incorporated herein by reference). The expression units may also include a transcriptional terminator. A preferred transcriptional terminator is the TPI1 terminator (Alber and Kawasaki, ibid.).

Additional vectors, promoters and terminators for use in expressing the transglutaminases of the invention in yeast are well known in the art and are reviewed by, for example, Emr, Meth. Enzymol. 185:231-279, (1990), incorporated
5 herein by reference.

The transglutaminases of the invention may be expressed in Aspergillus spp. (McKnight and Upshall, described in U.S. Patent 4,935,349, which is incorporated herein by reference). Useful promoters include those derived from
10 Aspergillus nidulans glycolytic genes, such as the ADH3 promoter (McKnight et al., EMBO J. 4:2093-2099, 1985) and the tpiA promoter. An example of a suitable terminator is the ADH3 terminator (McKnight et al., *ibid.*).

Techniques for transforming fungi are well known in
15 the literature, and have been described, for instance, by Beggs (*ibid.*), Hinnen et al. (Proc. Natl. Acad. Sci. USA 75: 1929-1933, 1978), Yelton et al. (Proc. Natl. Acad. Sci. USA 81: 1740-1747, 1984), and Russell (Nature 301: 167-169, 1983). The genotype of the host cell will generally contain a genetic
20 defect that is complemented by the selectable marker present on the expression vector. Choice of a particular host and selectable marker is well within the level of ordinary skill in the art.

In addition to fungal cells, cultured mammalian
25 cells may be used as host cells within the present invention. Preferred cultured mammalian cells for use in the present invention include the COS-1 (ATCC CRL 1650) and BALB/c 3T3 (ATCC CRL 163) cell lines. In addition, a number of other mammalian cell lines may be used within the present invention,
30 including BHK (ATCC CRL 10314), 293 (ATCC CRL 1573), Rat Hep I (ATCC CRL 1600), Rat Hep II (ATCC CRL 1548), TCMK (ATCC CRL 139), Human lung (ATCC CCL 75.1), Human hepatoma (ATCC HTB-52), Hep G2 (ATCC HB 8065), Mouse liver (ATCC CCL 29.1), NCTC 1469 (ATCC CCL 9.1) and DUKX cells (Urlaub and Chasin,
35 Proc. Natl. Acad. Sci. USA 77: 4216-4220, 1980).

Mammalian expression vectors for use in carrying out the present invention will include a promoter capable of directing the transcription of a cloned gene or cDNA. Preferred promoters include viral promoters and cellular promoters. Viral promoters include the immediate early cytomegalovirus promoter (Boshart et al., Cell 41: 521-530, 1985), the SV40 promoter (Subramani et al., Mol. Cell. Biol. 1: 854-864, 1981), and the major late promoter from Adenovirus 2 (Kaufman and Sharp, Mol. Cell. Biol. 2: 1304-1319, 1982). Cellular promoters include the mouse metallothionein-1 promoter (Palmiter et al., U.S. Patent No. 4,579,821), a mouse V_K promoter (Bergman et al., Proc. Natl. Acad. Sci. USA 81: 7041-7045, 1983; Grant et al., Nuc. Acids Res. 15: 5496, 1987) and a mouse V_H promoter (Loh et al., Cell 33: 85-93, 1983). Also contained in the expression vectors is a polyadenylation signal located downstream of the coding sequence of interest. Polyadenylation signals include the early or late polyadenylation signals from SV40 (Kaufman and Sharp, *ibid.*), the polyadenylation signal from the Adenovirus 5 E1B region and the human growth hormone gene terminator (DeNoto et al., Nuc. Acids Res. 9: 3719-3730, 1981). Vectors can also include enhancer sequences, such as the SV40 enhancer and the mouse μ enhancer (Gillies, Cell 33: 717-728, 1983). Expression vectors may also include sequences encoding the adenovirus VA RNAs. Vectors can be obtained from commercial sources (e.g., Stratagene, La Jolla, CA).

Cloned DNA sequences may be introduced into cultured mammalian cells by, for example, calcium phosphate-mediated transfection (Wigler et al., Cell 14: 725, 1978; Corsaro and Pearson, Somatic Cell Genetics 7: 603, 1981; Graham and Van der Eb, Virology 52: 456, 1973), electroporation (Neumann et al., EMBO J. 1: 841-845, 1982), DEAE-dextran mediated transfection (Ausubel et al., (ed.) Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY (1987), incorporated herein by reference) or a

commercially available transfection reagent and method such as the Boehringer Mannheim Transfection-Reagent N-[1-(2,3-Dioleoyloxy)propyl]-N,N,N-trimethyl ammoniummethylsulfate (Boehringer Mannheim, Indianapolis, IN). To identify cells
5 that have stably integrated the cloned DNA, a selectable marker is generally introduced into the cells along with the gene or cDNA of interest. Preferred selectable markers for use in cultured mammalian cells include genes that confer resistance to drugs, such as neomycin, hygromycin, and
10 methotrexate. The selectable marker may be an amplifiable selectable marker. Preferred amplifiable selectable markers are the DHFR gene and the neomycin resistance gene. Selectable markers are reviewed by Thilly (Mammalian Cell Technology, Butterworth Publishers, Stoneham, MA, which is
15 incorporated herein by reference). The choice of selectable markers is well within the level of ordinary skill in the art.

Selectable markers may be introduced into the cell on a separate vector at the same time as the transglutaminase sequence of interest, or they may be introduced on the same
20 vector. If on the same vector, the selectable marker and the transglutaminase sequence of interest may be under the control of different promoters or the same promoter, the latter arrangement producing a dicistronic message. Constructs of this type are known in the art (for example, Levinson and
25 Simonsen, U.S. Patent No. 4,713,339). It may also be advantageous to add additional DNA, known as "carrier DNA" to the mixture which is introduced into the cells.

Transfected mammalian cells are allowed to grow for a period of time, typically 1-2 days, to begin expressing the
30 DNA sequence(s) of interest. Drug selection is then applied to select for growth of cells that are expressing the selectable marker in a stable fashion. For cells that have been transfected with an amplifiable selectable marker the drug concentration may be increased in a stepwise manner to

select for increased copy number of the cloned sequences, thereby increasing expression levels.

Promoters, terminators and methods for introducing expression vectors encoding transglutaminase into plant, avian and insect cells are well known in the art. The use of baculoviruses, for example, as vectors for expressing heterologous DNA sequences in insect cells has been reviewed by Atkinson et al. (Pestic. Sci. 28: 215-224, 1990). The use of Agrobacterium rhizogenes as vectors for expressing genes in plant cells has been reviewed by Sinkar et al. (J. Biosci. (Bangalore) 11: 47-58, 1987).

Host cells containing DNA constructs of the present invention are then cultured to produce the transglutaminase. The cells are cultured according to standard methods in a culture medium containing nutrients required for growth of the chosen host cells. A variety of suitable media are known in the art and generally include a carbon source, a nitrogen source, essential amino acids, vitamins and minerals, as well as other components, e.g., growth factors or serum, that may be required by the particular host cells. The growth medium will generally select for cells containing the DNA construct by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker on the DNA construct or co-transfected with the DNA construct.

Yeast cells, for example, are preferably cultured in a medium which comprises a nitrogen source (e.g., yeast extract), inorganic salts, vitamins and trace elements. The pH of the medium is preferably maintained at a pH greater than 2 and less than 8, preferably at pH 5-6. Methods for maintaining a stable pH include buffering and constant pH control, preferably through the addition of sodium hydroxide. Preferred buffering agents include succinic acid and Bis-Tris (Sigma Chemical Co., St. Louis, MO). Cultured mammalian cells are generally cultured in commercially available serum-containing or serum-free media. Selection of a medium

appropriate for the particular cell line used is within the level of ordinary skill in the art.

In a preferred embodiment, the human prostate and placental transglutaminases are expressed in yeast as intracellular products. The yeast host can be a diploid strain homozygous for pep4, a mutation that reduces vacuolar protease levels, as described in Jones et al., Genetics 85:23-33 (1977), incorporated herein by reference. The strain is also homozygous for disruption of the endogenous TPI (triose phosphate isomerase) gene, thereby allowing the S. pombe POT1 gene to be used as a selectable marker. The vector includes the POT1 marker, a leu2-d marker and the ADH2-4^C promoter. The POT1 marker in the TPI⁻ host allows for selection by growth in glucose. The host strain is grown in glucose-containing synthetic media with a glucose feed. An ethanol feed is then substituted for glucose to de-repress the promoter. The pH is maintained with NaOH. Other preferred means for expression are generally described in, e.g., EPO publication EP 268,772, incorporated herein by reference.

In another preferred embodiment, the human prostate and placental transglutaminases are expressed in cultured mammalian cells. Preferably, the cultured mammalian cells are BHK 570 cells (deposited with the American Type Culture Collection under accession number 10314).

The human prostate and placental transglutaminases produced according to the present invention may be purified by affinity chromatography on an antibody column using antibodies directed against the transglutaminases. Additional purification may be achieved by conventional chemical purification means, such as liquid chromatography, gradient centrifugation, and gel electrophoresis, among others. Methods of protein purification are known in the art (see generally, Scopes, R., Protein Purification, Springer-Verlag, NY (1982), which is incorporated herein by reference) and may be applied to the purification of the recombinant

transglutaminase described herein. Antibodies prepared against the novel transglutaminases may be either polyclonal or monoclonal, and can be used to isolate and substantially purify the recombinant or native transglutaminases of the invention. Substantially pure recombinant human prostatic or placental transglutaminase of at least about 50% is preferred, at least about 70-80% more preferred, and 95-99% or more homogeneity most preferred, particularly for pharmaceutical uses. Once purified, partially or to homogeneity, as desired, the recombinant or native human prostatic and placental transglutaminases described herein may be used as desired.

The human prostatic and placental transglutaminases produced according to the present invention find a variety of uses. These transglutaminases can be used therapeutically in humans or other mammals. For example, human transglutaminase may be used in the repair of wounds, ulcerated lesions, skin grafts, etc. As the human transglutaminases are relatively stable, active extracellularly, and bind avidly to collagen, they can be used to stabilize basement membrane structures. An appropriate endogenous substrate for transglutaminase is fibronectin, which thus serves as a basis for crosslinking and stabilizing collagen/fibronectin complexes.

Pharmaceutical compositions of the invention comprise therapeutically effective amounts of human prostatic and/or placental transglutaminase and an appropriate physiologically acceptable carrier. The pharmaceutical compositions are intended primarily for topical or local administration, for use in methods of wound closure, as tissue adhesives, and the like. Typically the transglutaminase will be administered concurrently with or prior to compositions of thrombin to the wound site to increase effectiveness.

A variety of aqueous carriers may be used in the compositions, e.g., water, buff red wat r, saline, 0.3% glycine and the like, including glycoproteins for enhanced stability, such as albumin, lipoprotein, fibronectin and/or

globulin. The compositions may be sterilized by well known sterilization techniques, and the solutions packaged for use or lyophilized. Other components of the pharmaceutical compositions of the invention can include pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, etc.

Other components may also be added to the transglutaminase compositions to enhance their effectiveness, such as calcium ions, protease inhibitors (e.g., aprotinin), fibrinogen, etc. Admixtures of prostaglandins, coagulation factors, antihistamines, vasopressins, growth factors, vitamins, antibiotics (e.g., aminoglycosides, penicillins, carbapenems, sulfonamides, tetracyclines) and the like may also be provided. The formulation of various wound tissue adhesives is discussed in detail in U.S. Pat. Nos. 4,427,650, 4,442,655, and 4,655,211, each of which is incorporated herein by reference.

The concentration of human prostatic and/or placental transglutaminase in the pharmaceutical formulations can vary widely, i.e., from about 20 $\mu\text{g/ml}$ to 20 mg/ml or more, usually at least about 50 μg to 1 mg/ml , preferably from about 100 μg to 500 $\mu\text{g/ml}$ and will be selected primarily by volumes, viscosities, strength of the resulting complex, etc., in accordance with the particular use intended, the severity of the wound, the mode of administration selected, etc. Amounts effective for these uses will depend on the severity of the wound, injury or disease and the general state of the patient, but generally range from about 100 μg to about 500 mg or more of transglutaminase per site, with dosages of from about 500 μg to about 50 mg of transglutaminase per site being more commonly used. It must be kept in mind that the materials of the present invention may generally be employed

in serious disease or injury states, that is, life-threatening or potentially life threatening situations. In such cases, in view of the minimization of extraneous substances, decreased immunogenicity and the prolonged half-life and stability of the human prostatic and placental transglutaminases made feasible by this invention, it is possible and may be felt desirable by the treating physician to administer substantial excesses of these transglutaminase compositions.

The transglutaminases described herein can also be used in the preparation of food material, such as paste food or cheese, and can be added to dehydrated fish to prevent deterioration caused by protozoans, e.g., myxamoeba. The transglutaminases can also be used in the preparation of ground meat of okiomi (Euphasia superba), by adding to dehydrated meat parts from 0.1 to 100 units, preferably about 1-40 U per gram of protein to improve meat texture and quality. Frozen granular meats can be improved by combining meat material with transglutaminase of the invention at 1-500 U per gram protein, at 30-60°C for 10-120 min. to promote crosslinking between glutamine groups and lysine contained in meat preparations.

Other uses of the human prostatic and placental transglutaminases described herein include use in the enzyme-catalyzed labeling of proteins and cell membranes (Iwani, Eur. J. Biochem. 80:359-368 (1977), incorporated herein by reference), in the introduction of cleavable crosslinks, and in the solid phase reversible removal of specific proteins from biological systems.

Transglutaminase expression can be used as a marker for screening for agonists and antagonists of cellular apoptosis. Identifying agents which inhibit the expression of transglutaminase by a cell provides a means to prevent or delay atrophic changes characteristic of many degenerative changes, particularly degenerative nerve diseases, such as Parkinson's disease and Alzheimer's disease. Inhibition of

apoptosis may also enhance blood cell counts in chemotherapy patients. The human prostatic and placental transglutaminase or the nucleic acids which encode the transglutaminases of the invention can also be used to identify agents which induce apoptotic activity by a cell, for the control of, e.g., hyperproliferative disorders. The growth of cells such as adipocytes can be regulated with agents identified using the transglutaminases provided herein as a marker, providing a means for controlling fat deposits in certain forms of obesity without the necessity for surgical intervention.

Polynucleotide molecules which encode the prostatic and placental transglutaminases may be directly detected in cells with labeled synthetic oligonucleotide probes in a hybridization procedure similar to the Southern or dot blot. Also, PCR (including Saiki et al., Science 239:487 (1988)) may be used to amplify DNA sequences, which are subsequently detected by their characteristic size on agarose gels, Southern blots of the gels using transglutaminase sequences or oligonucleotide probes, or a dot blot using similar probes. The probes of the present invention are at least 85% homologous to a corresponding DNA sequence of a human prostate transglutaminase sequence of Sequence ID No. 14 or its complement or a human placental transglutaminase sequence of Sequence ID No. 22 or its complement. For use as probes, the molecules may comprise from about 14 nucleotides to about 25 or more nucleotides, sometimes 40 to 60 nucleotides, and in some instances a substantial portion or even the entire cDNA of a transglutaminase gene of the invention may be used. The probes are labeled to provide a detectable signal, such as an enzyme, biotin, a radionuclide, fluorophore, chemiluminescer, paramagnetic particle, etc.

The following examples are provided by way of illustration, not limitation.

EXAMPLE I

Cloning of Human Prostatic Transglutaminase

5 This Example describes the construction of oligonucleotide primers for amplification via PCR of sequences encoding human prostate transglutaminase, the cloning of the human prostate transglutaminase gene, and its nucleotide sequencing.

10 A series of synthetic degenerate oligonucleotide primers were generated to encode three regions of conserved amino acid sequences identified from a multiple alignment of known transglutaminase sequences, human erythrocyte membrane protein band 4.2 and the rat dorsal protein-1 (Ho et al., Prog. Clin. Biol. Res. 239: 125-153, (1987)). The multiple
15 alignment employed sequences of rat keratinocyte transglutaminase, human keratinocyte transglutaminase, human transglutaminase K, human factor XIII, human endothelial cell transglutaminase, mouse macrophage transglutaminase, guinea pig transglutaminase, human erythrocyte membrane protein band
20 4.2, rabbit transglutaminase type I, and bovine factor XIII. The multiple alignment was subjected to analysis of subsequence for least degenerate/most conserved regions to design primers of 17-20 bases in length. The amino acid sequences across three regions of multiple homology were
25 chosen as the basis from which to design degenerate primers: One region corresponding to the active site region of factor XIII, and two other regions which seemed to have structural importance, based on, inter alia, the presence of hydrophobic residues and proline residues. Degenerate oligonucleotides
30 ZC4109, ZC4110, ZC4111 and ZC4112 (Sequence ID Nos. 1, 2, 3 and 4) were designed to provide DNA segments corresponding to the conserved amino acid coding sequences. Degenerate oligonucleotides ZC4120, ZC4121, ZC4122, ZC4127, ZC4128, and
35 ZC4129 (Sequence ID Nos. 5, 6, 7, 8, 9 and 10; Table 1) were designed such that each primer contained a 5' prime sequence

to facilitate cloning into prime vectors described by Hagen (U.S. Patent No. 5,075,227, incorporated herein by reference) in addition to a DNA segment corresponding to the conserved amino acid coding sequence. The prime sequences shown in Table 1 are underlined.

Table 1: Degenerate Oligonucleotide Primers (5' to 3')

ZC4120 (Sequence ID Number 5)

CATCCACGGA CTACGACGAR TAYSTNCTNA MYGA

ZC4121 (Sequence ID Number 6)

CATCCACGGA CTACGACGAR TAYSTNCTNA MRGA

ZC4122 (Sequence ID Number 7)

CATCCACGGA CTACGACGAR TAYSTNCTNA MNCA

ZC4127 (Sequence ID Number 8)

CATCCACGGA CTACGACTAY GGNCARTGY TGGGTNTT

ZC4128 (Sequence ID Number 9)

ACTCTCCGGT ACGACAGAAN ACCCARCAYT GNCC

ZC4129 (Sequence ID Number 10)

ACTCTCCGGT ACGACAGCCY TCNKGRWRYT TRTA

The oligonucleotide primers were paired as shown in Table 2, and each pair was used in a PCR reaction using a λ gt11 human prostate tissue cDNA library obtained from Clontech Laboratories, Inc., Palo Alto, CA as a template. Fifty microliter reactions were set up with each reaction containing 0.2 mM each of dCTP, dGTP, dATP and dTTP, 2 pmol of each primer, 1 μ l of the cDNA library, 3 units of Taq polymerase (Promega Corp., Madison, WI) and 5 μ l of 10x Promega PCR buffer (Promega Corp., Madison, WI). The reactions were each overlaid with mineral oil and amplified

with two cycles (90 seconds at 94°C, 90 seconds at 40°C, 2 minutes at 72°C), thirty-eight cycles (45 seconds at 94°C, 45 seconds at 45°C, two minutes at 72°C) and an incubation at 72°C for seven minutes.

5

Table 2: Oligonucleotide Primer Combinations And Expected
Fragment Sizes (Base Pairs)

RXN	SENSE OLIGO	ANTISENSE OLIGO	EXPECTED FRAGMENT SIZE
10	1. ZC4110	ZC4112	344
	2. ZC4110	ZC4111	851
	3. ZC4109	ZC4111	527
	4. ZC4127	ZC4129	561
	5. ZC4120	ZC4128	378
	6. ZC4121	ZC4128	378
15	7. ZC4122	ZC4128	378
	8. ZC4120	ZC4129	885
	9. ZC4121	ZC4129	885
	10. ZC4122	ZC4129	885

20

Aliquots of the reaction mixtures were electrophoresed on an agarose gel. Reactions 3, 4 and 6 exhibited bands of expected size (527 bp, 561 bp and 378 bp, respectively). The PCR reaction products were isolated by agarose gel electrophoresis, and the DNA fragments were extracted with a Bio-Rad PREP-A-GENE kit (Bio-Rad, Richmond, CA) using the manufacturer's directions. The purified fragments were ligated into pCR1000 (Invitrogen, San Diego, CA) from the TA Cloning Kit (Invitrogen) and transformed into *E. coli* strain INVaF' (Invitrogen) using the manufacturer's protocol (Invitrogen TA Cloning Instruction Manual K2000-1). Two clones from PCR reaction 4, designated PTG561/1 and PTG561/2, were selected for subsequent analysis. Sequence analysis of the PCR-generated cDNA inserts in plasmids PTG561/1 and PTG561/2 showed that PTG561/2 (SEQ. ID. NO. 13) contained a unique sequence.

25

30

35

To generate a full-length prostate transglutaminase cDNA clone, sense and antisense oligonucleotide primers were designed to specific sequences in the PTG561/2 clone.

Oligonucleotides ZC4248 and ZC4249 (Sequence ID Nos. 11 and 12) were used to amplify a 468 base pair fragment from clone PTG561/2 that was used to probe the prostate cDNA library. A fifty microliter reaction mixture was set up containing 2 pmols each of ZC4248 and ZC4249, 0.025 mM dGTP, 0.025 mM dTTP, 6.6×10^{-3} mM α -³²P dCTP, 6.6×10^{-3} mM α -³²P dATP, 1x Promega PCR buffer, 1 μ l of purified plasmid diluted 1:100 and 0.5 μ l Taq polymerase. The reaction mixture was layered with mineral oil, and the mixture was preheated for approximately three minutes at 87°C. The reaction was amplified for six cycles (one minute at 94°C, one minute at 45°C, one minute at 72°C) and one incubation at 72°C for five minutes. A 45 μ l aliquot of the amplified reaction mixture was isopropanol precipitated, and the radiolabeled PCR product was used to probe the λ gt11 human prostate cDNA library (Clontech). Six positive clones were selected for further analysis.

The six clones were subjected to PCR amplification using oligonucleotides ZC4362 and ZC4363 (Sequence ID Nos. 19 and 20, respectively), which were designed to anneal to sequences in the λ gt11 vector, to characterize the cDNA inserts. Six 50 μ l reaction mixtures were prepared, each of which contained 1 μ l of a plate lysate of one of the selected clones, 1X Promega PCR buffer, 0.2 mM of each dNTP, 2 pmole each of ZC4362 and ZC4363 (Sequence ID Nos. 19 and 20, respectively), and 3 units of Taq polymerase. The reactions were overlaid with mineral oil, and the mixtures were pre-heated to 94°C for two minutes to disrupt the phage. The reactions were amplified through 30 cycles (1 minute at 94°C, 1 minute at 50°C, three minutes at 72°C) followed by one cycle at 72°C for 7 minutes. The reaction products were isolated by agarose gel electrophoresis, and each reaction product was subcloned into pCR1000 (Invitrogen, San Diego CA) and

transformed into *E. coli* strain INV α F' (Invitrogen) using the TA cloning kit (Invitrogen).

One of the six lambda clones, 8c2, was selected for sequence analysis. Lambda DNA was prepared from the 8c2 clone, and the cDNA insert was isolated as an Eco RI fragment and subcloned into Eco RI-linearized pUC18 to obtain plasmid pDT43. The 8c2 cDNA insert was subjected to DNA sequence analysis. Based on homology with a published rat prostate protein sequence (Ho et al., *J. Biol. Chem.* 267: 12660-12667, 1992), it was determined that the prostate transglutaminase clone 8c2 lacked the 5' coding sequence.

To confirm the presence of additional 5' sequences, the original six lambda clones were used as templates for PCR reactions using oligonucleotides ZC5509 (Sequence ID No. 21) and ZC4048 (Sequence ID No. 18). Each reaction mixture contained 1x PCR buffer, 1.25 mM MgCl₂, 0.2 mM of each dNTP, 20 μ M ZC5509, 17.5 μ l of 20 μ M ZC4048, 1.5 units of Taq polymerase. The reaction mixture was divided into 24 μ l aliquots. Each aliquot received 1 μ l of template, and the reaction mixtures were amplified for thirty cycles (94°C for one minute, 42°C for one minute, 72°C for two minutes) followed by a seven minute incubation at 72°C. The reaction mixtures were subjected to agarose gel electrophoresis. Analysis of the PCR products showed that clones 11A2 and 11A3 generated the largest PCR products relative to clone 8c2, suggesting that these two clones contained additional 5' prostate transglutaminase coding sequences.

The 5' human prostate coding sequence was obtained by amplification from the two lambda clones (11A2 and 11A3) described above. Synthetic oligonucleotide ZC4048 (Sequence ID No. 18) was designed to hybridize to the antisense lambda sequences near the Eco RI site of the λ gt11 vector. Synthetic oligonucleotide ZC5509 (Sequence ID No. 21) was designed to hybridize to the sense sequences in the 5' coding sequence of the PTG561/2 cDNA (Sequence ID No. 13).

25

Two 50 μ l reaction mixture were prepared containing 9.3 μ l of either 11A2 or 11A3 phage from plate lysates, 5 μ l 10X Promega PCR buffer, 5 μ l of a solution containing 0.2 mM of each dNTP, 2.5 μ l each of 20 pMol/ μ l ZC4048 and 20 pMol/ μ l ZC5509 (Sequence ID Nos. 18 and 21, respectively), 25.1 μ l of water and 0.6 μ l of Taq polymerase. The reactions were incubated at 94°C for two minutes to disrupt the phage followed by thirty cycles (45 seconds at 94°C, 45 seconds at 42°C, 90 seconds at 72°C). After the final amplification cycle, the reactions were incubated at 72°C for five minutes. The reactions were subjected to agarose gel electrophoresis, and an approximately 530 bp band was isolated from each reaction. The PCR-generated fragments were subcloned into pCRII (Invitrogen, San Diego CA) using the manufacturer's supplied instructions. Sequence analysis of several clones showed identical sequences spanning the λ gt11 Eco RI cloning site and sequences present in the 8c2 clone. One clone, pDT46-1 was selected for further manipulation.

The 5' transglutaminase coding sequence present in pDT46-1 but missing from the 8c2 clone was obtained by digesting pDT46-1 with Spe I and Ava I to isolate the 351 bp fragment. The 3' transglutaminase coding sequence was obtained by digesting plasmid pDT43 with Ava I and Xba I and isolating the fragment containing the transglutaminase and vector sequences. The Spe I and Xba I digestion produce complementary adhesive ends. The Spe I-Ava I fragment from pDT46-1 and the Ava I-Xba I fragment from pDT43 were ligated to obtain plasmid pDT47-15, which contained the prostate transglutaminase coding sequence of Sequence ID No. 14.

EXAMPLE II

Expression of Human Prostate Transglutaminase

This Example describes the expression of a human prostate transglutaminase from cultured mammalian cells.

The prostate transglutaminase cDNA insert present in plasmid pDT47-15 was subcloned into the mammalian expression vector Zem229R. Plasmid Zem229 is a pUC18-based expression vector containing a unique Bam HI site for insertion of cloned DNA between the mouse metallothionein-1 promoter and SV40 transcription terminator and an expression unit containing the SV40 early promoter, mouse dihydrofolate reductase gene, and SV40 terminator. Zem229 was modified to delete the two Eco RI sites by partial digestion with Eco RI, blunting with DNA polymerase I (Klenow fragment) and dNTPs, and re-ligation. Digestion of the resulting plasmid with Bam HI followed by ligation of the linearized plasmid with Bam HI-Eco RI adapters resulted in a unique Eco RI cloning site. The resultant plasmid was designated Zem229R.

Plasmid pDT47-15 was digested with Hind III to isolate the approximately 3 kb Hind III fragment containing the prostate transglutaminase cDNA. Synthetic oligonucleotides ZC1157 and ZC1158 (Sequence ID Nos. 16 and 17, respectively) were kinased and annealed to form Eco RI-Hind III adapters. The kinased, annealed oligonucleotides and the 3 kb Hind III fragment were ligated to Eco RI-linearized Zem229R. The ligation mixture was transformed into *E. coli* strain DH10B cells, and transformants were selected for growth in the presence of ampicillin. Plasmid DNA prepared from selected transformants was subjected to restriction endonuclease analysis. A plasmid clone, pPTG/229R, suspected of having the insert in the correct orientation relative to the promoter, was selected for DNA sequence analysis to confirm the orientation of the insert. DNA sequence analysis confirmed the orientation of the insert and also disclosed the presence of polylinker sequences between the promoter sequence of Zem229R and the beginning of the prostate transglutaminase coding sequence. These

sequences appeared to be remnants from the initial cloning procedure.

The polylinker sequences between the promoter sequence of Zem229R and the prostate transglutaminase coding sequence were removed by first digesting plasmid pPTG/229R with Eco RI to completion. The approximately 236 base pair fragment containing the 5'-most transglutaminase coding sequence and the approximately 2.7 kb fragment containing the remainder of the transglutaminase coding sequence were isolated by agarose gel electrophoresis. The two Eco RI fragments were ligated with Eco RI-linearized Zem229R that had been treated with calf alkaline phosphatase to prevent recircularization. The ligation mixture was transformed into *E. coli* strain DH10B cells, and transformants were selected in the presence of ampicillin. Plasmid DNA prepared from selected transformants was analyzed by restriction enzyme analysis. A plasmid containing the prostate transglutaminase cDNA insert in the proper orientation relative to the promoter in Zem229R was designated pPTGR/229R.

Both plasmids pPTG/229R and pPTGR/229R were transfected into BHK 570 cells (deposited with the American Type Culture Collection under accession number 10314) using Boehringer Mannheim Transfection-Reagent N-[1-(2,3-Dioleoyloxy)propyl]-N,N,N-trimethyl ammoniummethylsulfate using the manufacturer-supplied directions. The cells were cultured under non-selective conditions for two days. After two days the pPTG/229R transfectants were selected in media containing 1 μ M methotrexate, and the pPTGR/229R transfectants were selected in media containing either 1 μ M or 10 μ M methotrexate.

Transfectant colonies were overlaid with a nitrocellulose filter, and the colonies were incubated for 3 hours. After incubation, the filter was lifted and probed with rabbit anti-rat prostate transglutaminase antiserum, obtained from Dr. V. Gentile (University of Texas-Medical

School, Houston, Texas). The filters were incubated with a peroxidase-conjugated goat anti-rabbit IgG, and colonies bound by the rabbit anti-rat prostate transglutaminase antibodies were visualized using the chemiluminescent ECL REAGENT (Amersham Corp., Arlington Heights, IL) using the manufacturer's instructions. Six positive pPTG/229R transfectant colonies were each picked into a well of a 24-well plate. Of the pPTGR/229R transfectant colonies, 12 positive colonies were picked from those colonies selected in the presence of 1 μ M methotrexate, and 12 positive colonies were picked from those colonies selected in the presence of 10 μ M methotrexate.

The colonies are subjected to in vivo labeling followed by radioimmunoprecipitation of the protein with the rabbit anti-rat prostate transglutaminase antiserum. Briefly, the medium in each well is replaced with 1 ml of serum-free medium (DMEM -Lys -Met, 1 mM sodium pyruvate, 2mM L-glutamine, 5 mg/l insulin, 2 μ g/l selenium, 10 mg/l fetuin, 10 mg/l transferrin and 25 mM pH 7.2 HEPES buffer) containing 20 μ Ci of 35 S-EXPRESS (Du Pont-NEN Research Products, Boston MA), and the cells are incubated overnight at 37°C. After incubation, 1 ml of each supernatant is harvested, and the cells are rinsed with PBS. Cell extracts from each culture are obtained by incubating the cells with 1 ml RIPA buffer (10 mM Tris, pH 7.4, 1% deoxycholate, 1% Triton X-100, 0.1% SDS, 5 mM EDTA, 0.7 M NaCl). The labeled proteins are incubated with a 1:300 dilution of rabbit anti-rat prostate transglutaminase antiserum on ice for one hour. After incubation, 10 μ l of PANSORBIN (S. aureus cells, Calbiochem, San Diego, CA) is added to each reaction, and the mixtures are incubated on ice for one hour. The reactions are centrifuged, and the pellets are resuspended in 1 ml of RIP wash buffer (0.1% SDS, 5 mM EDTA, 0.7 M NaCl). The reactions are centrifuged, the pellets are each resuspended in 20 μ l of loading buffer and the samples are applied to a 10/20 gradient polyacrylamide gel

(Daichi). The gel is fixed for thirty minutes in 40% methanol, 10% acetic acid, following which the gel is incubated in AMPLIFY (Amersham) for 30 minutes. The gel is dried and exposed to film at -80°C. The autoradiograph is examined to identify the presence of prostate transglutaminase.

EXAMPLE III

10 Cloning of Human Placental Transglutaminase

This Example describes the cloning of a human placental transglutaminase from human placental cDNA and identification of a confirmatory human prostatic transglutaminase clone from human liver cDNA.

Two other cDNA sources were used in conjunction with selected degenerate oligonucleotide primers described in Example I to obtain unique transglutaminase cDNAs. QUICK-CLONE human liver cDNA (Clontech) and QUICK-CLONE human placenta cDNA (Clontech) were used as templates with oligonucleotide primers paired as shown in Table 3. Fifty microliter reactions were set up with each reaction containing 0.2 mM each of dCTP, dGTP, dATP and dTTP, 2 pmol of each primer, 1 µg of the cDNA library, 3 units of Taq polymerase (Promega Corp., Madison, WI) and 5 µl of 10x Promega PCR buffer (Promega Corp., Madison, WI). The reactions were layered with mineral oil and amplified with two cycles (90 seconds at 94°C, 90 seconds at 50°C, 2 minutes at 72°C), twenty-five cycles (45 seconds at 94°C, 45 seconds at 55°C, one minute at 72°C) and one incubation at 72°C for seven minutes.

Table 3: Oligonucleotide Primer Combinations And Expected
Fragment Sizes (Base Pairs)

RXN	TEMPLATE	SENSE OLIGO	ANTISENSE OLIGO	EXP. FRAG. SIZE
5	LIVER	ZC4127	ZC4129	561
		ZC4120	ZC4128	378
		ZC4121	ZC4128	378
		ZC4122	ZC4128	378
		ZC4120	ZC4129	885
10	PLACENTA	ZC4121	ZC4129	885
		ZC4122	ZC4129	885
		ZC4127	ZC4129	561
		ZC4120	ZC4128	378
		ZC4121	ZC4128	378
15		ZC4122	ZC4128	378
		ZC4120	ZC4129	885
		ZC4121	ZC4129	885
		ZC4122	ZC4129	885

20 Aliquots of the amplified DNA were electrophoresed on agarose gels. Reactions 1, 3, 14, 15, 16, and 17 yielded fragments of expected size (Table 3). The PCR-generated cDNA fragments were electrophoresed on agarose gels, and the fragments were extracted with a Bio-Rad PREP-A-GENE Kit (Bio-Rad, Richmond, CA) using the manufacturer's directions.

25 The purified fragments were ligated into pCR1000 (Invitrogen, San Diego, CA) and transformed into *E. coli* strain INVaF' (Invitrogen) according to the TA Cloning Kit (Invitrogen) using the manufacturer's protocol (Invitrogen TA Cloning Instruction Manual K2000-1). Clones from reactions 1 and 14 were selected for subsequent analysis. Sequence analysis of a clone arising from reaction 1 revealed the same human prostatic transglutaminase sequence as found in PTG561/2. Sequence analysis of a clone arising from reaction 14 PCR

35 cDNA, designated p1TG561/5, revealed a novel transglutaminase

sequence. The nucleotide sequence of p1TG561/5 is shown in Sequence ID No. 22.

It is evident from the above results that compositions are provided which encode novel prostatic and placental human transglutaminases. Pharmaceutical preparations of these transglutaminases are particularly useful as wound tissue adhesives, in view of the minimization of extraneous substances when produced by recombinant means, decreased immunogenicity in humans and prolonged half-life and stability. The efficacy, convenience of administration, and reduced cost are among the advantages conferred by the compositions of the invention.

The transglutaminases described herein can also be used, inter alia, in the preparation of food material, in the enzyme-catalyzed labeling of proteins and cell membranes, as markers for screening for agonists and antagonists of cellular apoptosis, and for the detection or monitoring of expression in cells with labeled synthetic oligonucleotide probes or other convenient assays.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: O'Hara, Patrick J
Grant, Francis J
Sheppard, Paul O
- (ii) TITLE OF INVENTION: NOVEL HUMAN TRANSGLUTAMINASES
- (iii) NUMBER OF SEQUENCES: 22
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Townsend and Townsend
 - (B) STREET: One Market Plaza, Steuart Street Tower
 - (C) CITY: San Francisco
 - (D) STATE: CA
 - (E) COUNTRY: USA
 - (F) ZIP: 94105-1492
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: US
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 07/816,284
 - (B) FILING DATE: 31-DEC-1991
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Parmelee, Steve W
 - (B) REGISTRATION NUMBER: 31-990
 - (C) REFERENCE/DOCKET NUMBER: 13952-13-1
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 206-467-9600
 - (B) TELEFAX: 206-623-6793

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZC4109
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TAYGGNCART GYTGGGTNTT

33

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZC4110

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GARTAYSTNC TNAMNSA

17

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZC4111

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCYTCNKGRW RYTTRTA

17

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZC4112

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

AANACCCARC AYTGNCC

17

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZC4120

34

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CATCCACGGA CTACGACGAR TAYSTNCTNA MYGA

34

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZC4121

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CATCCACGGA CTACGACGAR TAYSTNCTNA MRGA

34

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZC4122

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CATCCACGGA CTACGACGAR TAYSTNCTNA MNCA

34

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 37 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZC4127

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CATCCACGGA CTACGACTAY GGNCARTGYT GGGTNTT

37

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

35

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZC4128

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ACTCTCCGGT ACGACAGAA N ACCCARCAYT GNCC

34

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZC4129

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ACTCTCCGGT ACGACAGCCY TCNKGRWRYT TRTA

34

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZC4248

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

AGAGGCGATA TCTCTCCGCC TGTCTTGGCC CACTGC

36

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZC4249

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GGGATCCTGA CTACAGTGCT GAGAGCGTTG GGCATC

36

(2) INFORMATION FOR SEQ ID NO:13:

36

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 521 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 (B) CLONE: PTG562

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TATGGACAGT GCTGGGTATT TGCTGGGATC CTGACTACAG TGCTGAGAGC GTTGGGCATC	60
CCAGCACGCA GTGTGACAGG CTTGATTCA GCTCAGACA CAGAAAGGAA CCTCAGGTTG	120
GACACCTATG TGAATGAGAA TGGCGAGAAA ATCACCAGTA TGACCCACGA CTCTGTCTGG	180
AATTTCCATG TGTGGACGGA TGCCTGGATG AAGCGACCCT ACGACGGCTG GCAGGCTGTG	240
GACGCAACGC CGCAGGAGCG AAGCCAGGCT GTCTTCTGCT GTGGGCCATC ACCACTGACC	300
GCCATCCGCA AAGGTGACAT CTTTATTGTC TATGACACCA GATTCGTCTT CTCAGAAGTG	360
AATGGTGACA GGCTCATCTG GTTGGTGAAG ATGGTGAATG GGCAGGAGGA GTTACACGTA	420
ATTTCAATGG AGACCACAAG CATCGGGAAA AACATCAGCA CCAAGGCAGT GGGCCAAGAC	480
AGGCGGAGAG ATATCGCCTC TGAGTACAAG CTCCCCGAAG G	521

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 3064 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 147..2186

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AATTCTAAAA ATGCTTTTGC AAGCTTGCAAT GCCTGCAGGT GCAGCGGCCG CCAGTGTGAT	60
GGATATCTGC AGAATTCGGC TTGCGCTCAG CTGGAATTCC GCAGAGATAG AGTCTTCCCT	120
GGCATTGCAG GAGAGAATCT GAAGGG ATG ATG GAT GCA TCA AAA GAG CTG CAA	173
Met Met Asp Ala Ser Lys Glu Leu Gln	
1 5	
GTT CTC CAC ATT GAC TTC TTG AAT CAG GAC AAC GCC GTT TCT CAC CAC	221
Val Leu His Ile Asp Phe Leu Asn Gln Asp Asn Ala Val Ser His His	
10 15 20 25	
ACA TGG GAG TTC CAA ACG AGC AGT CCT GTG TTC CGG CGA GGA CAG GTG	269
Thr Trp Glu Phe Gln Thr Ser Ser Pro Val Phe Arg Arg Gly Gln Val	
30 35 40	

37

TTT	CAC	CTG	CGG	CTG	GTG	CTG	AAC	CAG	CCC	CTA	CAA	TCC	TAC	CAC	CAA	317
Phe	His	Leu	Arg	Leu	Val	Leu	Asn	Gln	Pro	Leu	Gln	Ser	Tyr	His	Gln	
			45					50					55			
CTG	AAA	CTG	GAA	TTC	AGC	ACA	GGG	CCG	AAT	CCT	AGC	ATC	GCC	AAA	CAC	365
Leu	Lys	Leu	Glu	Phe	Ser	Thr	Gly	Pro	Asn	Pro	Ser	Ile	Ala	Lys	His	
		60					65					70				
ACC	CTG	GTG	GTG	CTC	GAC	CCG	AGG	ACG	CCC	TCA	GAC	CAC	TAC	AAC	TGG	413
Thr	Leu	Val	Val	Leu	Asp	Pro	Arg	Thr	Pro	Ser	Asp	His	Tyr	Asn	Trp	
	75					80					85					
CAG	GCA	ACC	CTT	CAA	AAT	GAG	TCT	GGC	AAA	GAG	GTC	ACA	GTG	GCT	GTC	461
Gln	Ala	Thr	Leu	Gln	Asn	Glu	Ser	Gly	Lys	Glu	Val	Thr	Val	Ala	Val	
90					95					100					105	
ACC	AGT	TCC	CCC	AAT	GCC	ATC	CTG	GGC	AAG	TAC	CAA	CTA	AAC	GTG	AAA	509
Thr	Ser	Ser	Pro	Asn	Ala	Ile	Leu	Gly	Lys	Tyr	Gln	Leu	Asn	Val	Lys	
				110					115					120		
ACT	GGA	AAC	CAC	ATC	CTT	AAG	TCT	GAA	GAA	AAC	ATC	CTA	TAC	CTT	CTC	557
Thr	Gly	Asn	His	Ile	Leu	Lys	Ser	Glu	Glu	Asn	Ile	Leu	Tyr	Leu	Leu	
			125					130					135			
TTC	AAC	CCA	TGG	TGT	AAA	GAG	GAC	ATG	GTT	TTC	ATG	CCT	GAT	GAG	GAC	605
Phe	Asn	Pro	Trp	Cys	Lys	Glu	Asp	Met	Val	Phe	Met	Pro	Asp	Glu	Asp	
		140					145					150				
GAG	CGC	AAA	GAG	TAC	ATC	CTC	AAT	GAC	ACG	GGC	TGC	CAT	TAC	GTG	GGG	653
Glu	Arg	Lys	Glu	Tyr	Ile	Leu	Asn	Asp	Thr	Gly	Cys	His	Tyr	Val	Gly	
	155					160					165					
GCT	GCC	AGA	AGT	ATC	AAA	TGC	AAA	CCC	TGG	AAC	TTT	GGT	CAG	TTT	GAG	701
Ala	Ala	Arg	Ser	Ile	Lys	Cys	Lys	Pro	Trp	Asn	Phe	Gly	Gln	Phe	Glu	
170					175					180					185	
AAA	AAT	GTC	CTG	GAC	TGC	TGC	ATT	TCC	CTG	CTG	ACT	GAG	AGC	TCC	CTC	749
Lys	Asn	Val	Leu	Asp	Cys	Cys	Ile	Ser	Leu	Leu	Thr	Glu	Ser	Ser	Leu	
				190					195					200		
AAG	CCC	ACA	GAT	AGG	AGG	GAC	CCC	GTG	CTG	GTG	TGC	AGG	GCC	ATG	TGT	797
Lys	Pro	Thr	Asp	Arg	Arg	Asp	Pro	Val	Leu	Val	Cys	Arg	Ala	Met	Cys	
			205					210					215			
GCT	ATG	ATG	AGC	TTT	GAG	AAA	GGC	CAG	GGC	GTG	CTC	ATT	GGG	AAT	TGG	845
Ala	Met	Met	Ser	Phe	Glu	Lys	Gly	Gln	Gly	Val	Leu	Ile	Gly	Asn	Trp	
		220					225					230				
ACT	GGG	GAC	TAC	GAA	GGT	GGC	ACA	GCC	CCA	TAC	AAG	TGG	ACA	GGC	AGT	893
Thr	Gly	Asp	Tyr	Glu	Gly	Gly	Thr	Ala	Pro	Tyr	Lys	Trp	Thr	Gly	Ser	
	235					240					245					
GCC	CCG	ATC	CTG	CAG	CAG	TAC	TAC	AAC	ACG	AAG	CAG	GCT	GTG	TGC	TTT	941
Ala	Pro	Ile	Leu	Gln	Gln	Tyr	Tyr	Asn	Thr	Lys	Gln	Ala	Val	Cys	Phe	
250					255					260					265	
GGC	CAG	TGC	TGG	GTG	TTT	GCT	GGG	ATC	CTG	ACT	ACA	GTG	CTG	AGA	GCG	989
Gly	Gln	Cys	Trp	Val	Phe	Ala	Gly	Ile	Leu	Thr	Thr	Val	Leu	Arg	Ala	
				270					275					280		
TTG	GGC	ATC	CCA	GCA	CGC	AGT	GTG	ACA	GGC	TTC	GAT	TCA	GCT	CAC	GAC	1037
Leu	Gly	Ile	Pro	Ala	Arg	Ser	Val	Thr	Gly	Phe	Asp	Ser	Ala	His	Asp	
			285					290					295			

38

ACA Thr	GAA Glu	AGG Arg	AAC Asn	CTC Leu	ACG Thr	GTG Val	GAC Asp	ACC Thr	TAT Tyr	GTG Val	AAT Asn	GAG Glu	AAT Asn	GGC Gly	GAG Glu	1085
		300					305					310				
AAA Lys	ATC Ile	ACC Thr	AGT Ser	ATG Met	ACC Thr	CAC His	GAC Asp	TCT Ser	GTC Val	TGG Trp	AAT Asn	TTC Phe	CAT His	GTG Val	TGG Trp	1133
	315					320					325					
ACG Thr	GAT Asp	GCC Ala	TGG Trp	ATG Met	AAG Lys	CGA Arg	CCC Pro	TAC Tyr	GAC Asp	GGC Gly	TGG Trp	CAG Gln	GCT Ala	GTG Val	GAC Asp	1181
	330				335					340					345	
GCA Ala	ACG Thr	CCG Pro	CAG Gln	GAG Glu	CGA Arg	AGC Ser	CAG Gln	GGT Gly	GTC Val	TTC Phe	TGC Cys	TGT Cys	GGG Gly	CCA Pro	TCA Ser	1229
				350					355					360		
CCA Pro	CTG Leu	ACC Thr	GCC Ala	ATC Ile	CGC Arg	AAA Lys	GGT Gly	GAC Asp	ATC Ile	TTT Phe	ATT Ile	GTC Val	TAT Tyr	GAC Asp	ACC Thr	1277
			365					370					375			
AGA Arg	TTC Phe	GTC Val	TTC Phe	TCA Ser	GAA Glu	GTG Val	AAT Asn	GGT Gly	GAC Asp	AGG Arg	CTC Leu	ATC Ile	TGG Trp	TTG Leu	GTG Val	1325
		380					385					390				
AAG Lys	ATG Met	GTG Val	AAT Asn	GGG Gly	CAG Gln	GAG Glu	GAG Glu	TTA Leu	CAC His	GTA Val	ATT Ile	TCA Ser	ATG Met	GAG Glu	ACC Thr	1373
	395					400					405					
ACA Thr	AGC Ser	ATC Ile	GGG Gly	AAA Lys	AAC Asn	ATC Ile	AGC Ser	ACC Thr	AAG Lys	GCA Ala	GTG Val	GGC Gly	CAA Gln	GAC Asp	AGG Arg	1421
	410				415					420					425	
CGG Arg	AGA Arg	GAT Asp	ATC Ile	ACC Thr	TAT Tyr	GAG Glu	TAC Tyr	AAG Lys	TAT Tyr	CCA Pro	GAA Glu	GGC Gly	TCC Ser	TCT Ser	GAG Glu	1469
				430					435					440		
GAG Glu	AGG Arg	CAG Gln	GTC Val	ATG Met	GAT Asp	CAT His	GCC Ala	TTC Phe	CTC Leu	CTT Leu	CTC Leu	AGT Ser	TCT Ser	GAG Glu	AGG Arg	1517
			445					450					455			
GAG Glu	CAC His	AGA Arg	CAG Gln	CCT Pro	GTA Val	AAA Lys	GAG Glu	AAC Asn	TTT Phe	CTT Leu	CAC His	ATG Met	TCG Ser	GTA Val	CAA Gln	1565
		460					465					470				
TCA Ser	GAT Asp	GAT Asp	GTG Val	CTG Leu	CTG Leu	GGA Gly	AAC Asn	TCT Ser	GTT Val	AAT Asn	TTC Phe	ACC Thr	GTG Val	ATT Ile	CTT Leu	1613
	475					480					485					
AAA Lys	AGG Arg	AAG Lys	ACC Thr	GCT Ala	GCC Ala	CTA Leu	CAG Gln	AAT Asn	GTC Val	AAC Asn	ATC Ile	TTG Leu	GGC Gly	TCC Ser	TTT Phe	1661
	490				495				500						505	
GAA Glu	CTA Leu	CAG Gln	TTG Leu	TAC Tyr	ACT Thr	GGC Gly	AAG Lys	AAG Lys	ATG Met	GCA Ala	AAA Lys	CTG Leu	TGT Cys	GAC Asp	CTC Leu	1709
				510					515					520		
AAT Asn	AAG Lys	ACC Thr	TCG Ser	CAG Gln	ATC Ile	CAA Gln	GGT Gly	CAA Gln	GTA Val	TCA Ser	GAA Glu	GTG Val	ACT Thr	CTG Leu	ACC Thr	1757
		525						530					535			
TTG Leu	GAC Asp	TCC Ser	AAG Lys	ACC Thr	TAC Tyr	ATC Ile	AAC Asn	AGC Ser	CTG Leu	GCT Ala	ATA Ile	TTA Leu	GAT Asp	GAT Asp	GAG Glu	1805
		540					545					550				

39

CCA GTT ATC AGA GGT TTC ATC ATT GCG GAA ATT GTG GAG TCT AAG GAA Pro Val Ile Arg Gly Phe Ile Ile Ala Glu Ile Val Glu Ser Lys Glu 555 560 565	1853
ATC ATG GCC TCT GAA GTA TTC ACG TCA AAC CAG TAC CCT GAG TTC TCT Ile Met Ala Ser Glu Val Phe Thr Ser Asn Gln Tyr Pro Glu Phe Ser 570 575 580 585	1901
ATA GAG TTG CCT AAC ACA GGC AGA ATT GGC CAG CTA CTT GTC TGC AAT Ile Glu Leu Pro Asn Thr Gly Arg Ile Gly Gln Leu Leu Val Cys Asn 590 595 600	1949
TGT ATC TTC AAG AAT ACC CTG GCC ATC CCT TTG ACT GAC GTC AAG TTC Cys Ile Phe Lys Asn Thr Leu Ala Ile Pro Leu Thr Asp Val Lys Phe 605 610 615	1997
TCT TTG GAA AGC CTG GGC ATC TCC TCA CTA CAG ACC TCT GAC CAT GGG Ser Leu Glu Ser Leu Gly Ile Ser Ser Leu Gln Thr Ser Asp His Gly 620 625 630	2045
ACG GTG CAG CCT GGT GAG ACC ATC CAA TCC CAA ATA AAA TGC ACC CCA Thr Val Gln Pro Gly Glu Thr Ile Gln Ser Gln Ile Lys Cys Thr Pro 635 640 645	2093
ATA AAA ACT GGA CCC AAG AAA TTT ATC GTC AAG TTA AGT TCC AAA CAA Ile Lys Thr Gly Pro Lys Lys Phe Ile Val Lys Leu Ser Ser Lys Gln 650 655 660 665	2141
GTG AAA GAG ATT AAT GCT CAG AAG ATT GTT CTC ATC ACC AAG TAGCCTTGTC Val Lys Glu Ile Asn Ala Gln Lys Ile Val Leu Ile Thr Lys 670 675 680	2193
TGATGCTGTG GAGCCTTAGT TGAGATTTC A GCATTTCCTA CCTTGTGCTT AGCTTTTCAGA	2253
TTATGGATGA TTAAATTTGA TGACTTATAT GAGGGCAGAT TCAAGAGCCA GCAGGTCAAA	2313
AAGGCCAACA CAACCATAAG CAGCCAGACC CACAAGGCCA GGTCTGTGTC TATCACAGGG	2373
TCACCTCTTT TACAGTTAGA AACACCAGCC GAGGCCACAG AATCCCATCC CTTTCCTGAG	2433
TCATGGCCTC AAAAATCAGG GCCACCATTG TCTCAATTCA AATCCATAGA TTTTGAAGCC	2493
ACAGAGCTCT TCCCTGGAGC AGCAGACTAT GGGCAGCCCA GTGCTGCCAC CTGCTGACGA	2553
CCCTTGAGAA GCTGCCATAT CTTCAGGCCA TGGGTTTACC AGCCCTGAAG GCACCTGTCA	2613
ACTGGAGTGC TCTCTCAGCA CTGGGATGGG CCTGATAGAA GTGCATTCTC CTCCTATTGC	2673
CTCCATTCTC CTCTCTCTAT CCCTGAAATC CAGGAAGTCC CTCTCCTGGT GCTCCAAGCA	2733
GTTTGAAGCC CAATCTGCAA GGACATTTCT CAAGGGCCAT GTGGTTTTGC AGACAACCCT	2793
GTCCTCAGGC CTGAACTCAC CATAGAGACC CATGTCAGCA AACGGTGACC AGCAAATCCT	2853
CTTCCCTTAT TCTAAAGCTG CCCCTTGGGA GACTCCAGGG AGAAGGCATT GCTTCCTCCC	2913
TGGTGTGAAC TCTTTCTTTG GTATTCCATC CACTATCCTG GCAACTCAAG GCTGCTTCTG	2973
TTAACTGAAG CCTGCTCCTT CTGTGTTCTGC CCTCCAGAGA TTTGCTCAAA TGATCAATAA	3033
GCTTTAAATT AAACCGGAAT CCGCGGAATT C	3064

40

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 679 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

```

Met Met Asp Ala Ser Lys Glu Leu Gln Val Leu His Ile Asp Phe Leu
 1           5           10           15
Asn Gln Asp Asn Ala Val Ser His His Thr Trp Glu Phe Gln Thr Ser
 20           25           30
Ser Pro Val Phe Arg Arg Gly Gln Val Phe His Leu Arg Leu Val Leu
 35           40           45
Asn Gln Pro Leu Gln Ser Tyr His Gln Leu Lys Leu Glu Phe Ser Thr
 50           55           60
Gly Pro Asn Pro Ser Ile Ala Lys His Thr Leu Val Val Leu Asp Pro
 65           70           75           80
Arg Thr Pro Ser Asp His Tyr Asn Trp Gln Ala Thr Leu Gln Asn Glu
 85           90           95
Ser Gly Lys Glu Val Thr Val Ala Val Thr Ser Ser Pro Asn Ala Ile
100           105           110
Leu Gly Lys Tyr Gln Leu Asn Val Lys Thr Gly Asn His Ile Leu Lys
115           120           125
Ser Glu Glu Asn Ile Leu Tyr Leu Leu Phe Asn Pro Trp Cys Lys Glu
130           135           140
Asp Met Val Phe Met Pro Asp Glu Asp Glu Arg Lys Glu Tyr Ile Leu
145           150           155           160
Asn Asp Thr Gly Cys His Tyr Val Gly Ala Ala Arg Ser Ile Lys Cys
165           170           175
Lys Pro Trp Asn Phe Gly Gln Phe Glu Lys Asn Val Leu Asp Cys Cys
180           185           190
Ile Ser Leu Leu Thr Glu Ser Ser Leu Lys Pro Thr Asp Arg Arg Asp
195           200           205
Pro Val Leu Val Cys Arg Ala Met Cys Ala Met Met Ser Phe Glu Lys
210           215           220
Gly Gln Gly Val Leu Ile Gly Asn Trp Thr Gly Asp Tyr Glu Gly Gly
225           230           235           240
Thr Ala Pro Tyr Lys Trp Thr Gly Ser Ala Pro Ile Leu Gln Gln Tyr
245           250           255
Tyr Asn Thr Lys Gln Ala Val Cys Phe Gly Gln Cys Trp Val Phe Ala
260           265           270

```

41

Gly Ile Leu Thr Thr Val Leu Arg Ala Leu Gly Ile Pro Ala Arg Ser
 275 280 285
 Val Thr Gly Phe Asp Ser Ala His Asp Thr Glu Arg Asn Leu Thr Val
 290 295 300
 Asp Thr Tyr Val Asn Glu Asn Gly Glu Lys Ile Thr Ser Met Thr His
 305 310 315 320
 Asp Ser Val Trp Asn Phe His Val Trp Thr Asp Ala Trp Met Lys Arg
 325 330 335
 Pro Tyr Asp Gly Trp Gln Ala Val Asp Ala Thr Pro Gln Glu Arg Ser
 340 345 350
 Gln Gly Val Phe Cys Cys Gly Pro Ser Pro Leu Thr Ala Ile Arg Lys
 355 360 365
 Gly Asp Ile Phe Ile Val Tyr Asp Thr Arg Phe Val Phe Ser Glu Val
 370 375 380
 Asn Gly Asp Arg Leu Ile Trp Leu Val Lys Met Val Asn Gly Gln Glu
 385 390 395 400
 Glu Leu His Val Ile Ser Met Glu Thr Thr Ser Ile Gly Lys Asn Ile
 405 410 415
 Ser Thr Lys Ala Val Gly Gln Asp Arg Arg Arg Asp Ile Thr Tyr Glu
 420 425 430
 Tyr Lys Tyr Pro Glu Gly Ser Ser Glu Glu Arg Gln Val Met Asp His
 435 440 445
 Ala Phe Leu Leu Leu Ser Ser Glu Arg Glu His Arg Gln Pro Val Lys
 450 455 460
 Glu Asn Phe Leu His Met Ser Val Gln Ser Asp Asp Val Leu Leu Gly
 465 470 475 480
 Asn Ser Val Asn Phe Thr Val Ile Leu Lys Arg Lys Thr Ala Ala Leu
 485 490 495
 Gln Asn Val Asn Ile Leu Gly Ser Phe Glu Leu Gln Leu Tyr Thr Gly
 500 505 510
 Lys Lys Met Ala Lys Leu Cys Asp Leu Asn Lys Thr Ser Gln Ile Gln
 515 520 525
 Gly Gln Val Ser Glu Val Thr Leu Thr Leu Asp Ser Lys Thr Tyr Ile
 530 535 540
 Asn Ser Leu Ala Ile Leu Asp Asp Glu Pro Val Ile Arg Gly Phe Ile
 545 550 555 560
 Ile Ala Glu Ile Val Glu Ser Lys Glu Ile Met Ala Ser Glu Val Phe
 565 570 575
 Thr Ser Asn Gln Tyr Pro Glu Phe Ser Ile Glu Leu Pro Asn Thr Gly
 580 585 590
 Arg Ile Gly Gln Leu Leu Val Cys Asn Cys Ile Phe Lys Asn Thr Leu
 595 600 605

42

Ala Ile Pro Leu Thr Asp Val Lys Phe Ser Leu Glu Ser Leu Gly Ile
 610 615 620

Ser Ser Leu Gln Thr Ser Asp His Gly Thr Val Gln Pro Gly Glu Thr
 625 630 635 640

Ile Gln Ser Gln Ile Lys Cys Thr Pro Ile Lys Thr Gly Pro Lys Lys
 645 650 655

Phe Ile Val Lys Leu Ser Ser Lys Gln Val Lys Glu Ile Asn Ala Gln
 660 665 670

Lys Ile Val Leu Ile Thr Lys
 675

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
 (B) CLONE: ZC1157

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

AATTCTAAAA ATGCTTTTGC A

21

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
 (B) CLONE: ZC1158

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

AGCTTGCAAA AGCATTTTTA G

21

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
 (B) CLONE: ZC4048

43

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GCGCTCAGCT GGAAT

15

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZC4362

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

TGAATATCGA CGGTTTCCAT ATGG

24

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZC4363

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

TATTTTGGAC ACCAGACCAA CTGG

24

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZC5509

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

ACCATGTCCT CTTTACACCA T

21

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 527 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: pltG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TATGGTCAGT GTTGGGTTTT TGCTGGGACC CTCAACACAG CGCTGCGGTC TTTGGGGATT	60
CCTTCCCGGG TGATCACCAA CTTCAACTCA GCTCATGACA CAGACCGAAA TCTCAGTGTG	120
GATGTGTACT ACGACCCCAT GGGAAACCCC CTGGACAAGG GTAGTGATAG CGTATGGAAT	180
TTCCATGTCT GGAATGAAGG CTGGTTTGTG AGGTCTGACC TGGGCCCCCTC GTACGGTGGA	240
TGGCAGGTGT TGGATGCTAC CCCGCAGGAA AGAAGCCAAG GGGTGTTCCA GTGCGGCCCC	300
GCITCGGTCA TTGGTGTTTG AGAGGGTGAT GTGCAGCTGA ACTTCGACAT GCCCTTTATC	360
TTCGCGGAGG TTAATGCCGA CCGCATCACC TGGCTGTACG ACAACACCAC TGGCAAACAG	420
TGGAAGAATT CCGTGAACAG TCACACCATT GGCAGGTACA TCAGCACCAA GGCGGTGGGC	480
AGCAATGCTC GCATGGACGT CACGGACAAG TACAAGCTCC ACGAGGG	527

WHAT IS CLAIMED IS:

1. An isolated polynucleotide molecule which codes for human prostatic transglutaminase.

2. The polynucleotide molecule of claim 1, wherein the polypeptide encoded thereby catalyzes the Ca^{++} -dependent crosslinking of protein-bound glutamine and lysine residues.

3. The polynucleotide of claim 1, wherein the molecule is substantially the nucleotide sequence of human prostatic transglutaminase of Seq. ID. No. 14.

4. The polynucleotide of claim 1, which is a cDNA molecule.

5. An isolated polynucleotide molecule which codes for human placental transglutaminase.

6. The polynucleotide molecule of claim 5, wherein the polypeptide encoded thereby catalyzes the Ca^{++} -dependent crosslinking of protein-bound glutamine and lysine residues.

7. A DNA construct for the expression of human prostatic transglutaminase, which comprises the following operably linked elements:

a transcriptional promoter;
a DNA molecule encoding a human prostatic transglutaminase polypeptide; and
a transcriptional terminator.

8. A cultured cell transformed or transfected with the DNA construct of claim 7.

9. The cultured cell of claim 8, which is a eukaryotic cell.

5 10. The eukaryotic cell of claim 9, which is a yeast cell or mammalian cell.

10 11. A method for producing human prostatic transglutaminase, which comprises cultivating eukaryotic cells transformed or transfected with the DNA construct of claim 7, and isolating the transglutaminase from the cells.

12. The method of claim 11, wherein the transformed eukaryotic cells are yeast cells.

15 13. The polypeptide which is produced by the method of claim 11.

20 14. The polypeptide of claim 13, which catalyzes Ca^{++} -dependent crosslinking of protein-bound glutamine and lysine residues.

25 15. The polypeptide of claim 13, which has substantially the amino acid sequence of human prostatic transglutaminase of Seq. ID. No. 14.

30 16. A probe which comprises an oligonucleotide of at least about 14 nucleotides capable of specifically hybridizing with a gene which encodes a human prostatic or placental transglutaminase polypeptide, wherein said probe is at least 85% homologous to a sequence of the human prostatic or placental transglutaminase of Seq. ID. No. 14 or Seq. ID. No. 22 or its complement.

35 17. The probe of claim 16, which is labeled to provide a detectable signal.

18. A pharmaceutical composition which comprises purified human prostatic or placental transglutaminase and a pharmaceutically acceptable carrier.

5 19. A method for facilitating wound repair in a patient, which comprises administering the pharmaceutical composition of claim 19.

10 20. Purified recombinant human prostatic or placental transglutaminase.

15 21. An isolated polynucleotide molecule which codes for human placental transglutaminase, wherein the molecule comprises the nucleotide sequence of human placental transglutaminase of Seq. ID No. 22.

22. A DNA construct for the expression of human placental transglutaminase, which comprises the following operably linked elements:

20 a transcriptional promoter;
a DNA molecule encoding a human placental transglutaminase polypeptide; and
a transcriptional terminator.

25 23. A cultured cell transformed or transfected with the DNA construct of claim 22.

24. The transfected cell of claim 23, which is a yeast cell or mammalian cell.

30 25. A method for producing human placental transglutaminase, which comprises cultivating eukaryotic cells transformed or transfected with the DNA construct of claim 22, and isolating the transglutaminase from the cells.

26. The polypeptide which is produced by the method of claim 25.

5 27. The polypeptide of claim 26, which catalyzes Ca^{++} -dependent crosslinking of protein-bound glutamine and lysine residues.